

Supplementary file

Table S1. Primer sequences used for semi-quantitative reverse transcription and polymerase chain reaction (RT-PCR) analysis.

	Forward Primer (5'-3')	Reverse Primer (5'-3')
NF- κ B	AACAACACAGACCCAGGAGT	CTGTCACCAGGCGAGTTATAG
CCL2	GTCACCAAGCTCAAGAGAGAGA	GAGTGGATGCATTAGCTTCAGA
TNF- α	CGAGTCTGGGCAGGTCTACTTT	AGAGGTTGAGGGTGTCTGAAGG
IL-1 β	CCCTGAACTCAACTGTGAAATAGCA	CCCAAGTCAAGGGCTTGGA
IL-6	GAGTTGTGCAATGGCAATTC	ACTCCAGAAGACCAGAGCAG
AMPK	CGGCAAAGTGAAGGTTGGCAAA	CAAATAGCTCTCCTCCTGAGAC
Nrf2	TCTGACTCCGGCATTTCCT	GGCACTGTCTAGCTCTTCCA
OH-1	CACGCATATACCCGCTACCT	CCAGAGTGTTTCATTTCGAGA
β -actin	AAGTCCCTCACCTCCCAAAAG	AAGCAATGCTGTCACCTTCCC